

Anisotropic Local Stress and Particle Hopping in a Deeply Supercooled Liquid

Sarika Bhattacharyya and Biman Bagchi*

Solid State and Structural Chemistry Unit, Indian Institute of Science, Bangalore 560012, India

The origin of the microscopic motions that lead to stress relaxation in deeply supercooled liquid remains unclear. We show that in such a liquid the stress relaxation is locally anisotropic which can serve as the driving force for the hopping of the system on its free energy surface. However, not all hopping are equally effective in relaxing the local stress, suggesting that diffusion can decouple from viscosity even at local level. On the other hand, orientational relaxation is found to be always coupled to stress relaxation.

Dynamics of supercooled liquid show many fascinating properties, namely the non-exponentiality in various relaxation functions of dynamical variables, like the stress, density, composition, polarization and orientation [1, 2, 3, 4, 5]. One often observes a very slow power law decay in the intermediate to long times [6]. Although these have drawn attention of experimentalists and theoreticians and have been extensively studied, many aspects still remain ill-understood. It is believed that the nature of the microscopic motion undergoes a drastic change at a temperature T_B (a temperature substantially below the melting temperature, T_m). At the temperature T_B , the continuous hydrodynamic type motion, which is prevalent at higher temperature changes to discontinuous motion. It is also believed that T_B is close to the temperature where effects of the underlying free energy landscape on the dynamics are felt for the first time [1]. T_B is also found to be close to the mode coupling temperature, T_c . This change in the nature of the microscopic motion is believed to be the origin of the experimentally observed α - β bifurcation [7] and also the cross-over between the rotational and translational relaxation times [8].

Due to the complexity of the problem, computer simulations have played a key role in augmenting our understanding in this area. In particular, simulations allow one to directly look at the microscopic events. The computer simulation studies of the stress auto correlation function in the supercooled liquid could successfully reproduce the power law behavior of the stress auto-correlation function (SAF) [3, 4]. However, in the deeply supercooled liquid, one finds that within the simulation time, the relaxation, after an initial decay (typically 10-20%) becomes fully arrested. The microscopic origin of the subsequent decay is unclear. The computer simulation studies further show that the orientational and translational hopping of particles are the only mode present and hence the stress relaxation can happen only via hopping. However, since the relaxation time is much much longer (could be of the order of ms or sec), the computer simulation results, (which can explore mostly upto nano second regime and sometimes micro second regime), cannot include the ef-

fects of these hopping. Therefore, one cannot explore the relationship between hopping and the *total* stress relaxation.

The experiments on the other hand are successful in showing the decay of the SAF. However, except a recent work using single molecule spectroscopy [9], these experiments are macroscopic and do not provide enough information of the microscopic motions in the system.

Majumdar [10] had earlier discussed the importance of local relaxation modes (of wavelength less than the short range order) in giving rise to non-exponentiality in the stress relaxation function. This work discussed relaxation in terms of relaxation within small regions, surfaces and also volumes, with progressive lengthening of time scales. However, in that analysis the basic mechanism of relaxation was still assumed to be continuous.

In this Letter we demonstrate for the first time that in the deeply supercooled liquid (where hopping is the only surviving large amplitude motion), there is a close relationship between the *local* stress and the orientational and translational hopping. The local SAF is anisotropic and is found to change drastically during the hopping, thus showing that the local stress and the hopping of a particle are intimately connected. The anisotropy in the local stress could be the driving force for hopping. As the free energy of the system can be expressed in terms of the position dependent stress in a generalized Ginzburg-Landau formulation [11], the change of the anisotropic stress due to hopping should be regarded as the driving force for the transitions of the system between different minima of the free energy surface. However, not all hoppings are effective in relaxing the stress.

Our solvent is represented by binary Lennard-Jones mixture, which has been extensively studied [1, 12, 13, 14] and is known to be a good glass former, and our solute probes are prolate ellipsoids. Pressure is kept constant by Andersen's piston method [15] while in the case of temperature, a damped oscillator method has been adopted which keeps temperature constant at each and every time step [16]. The piston mass involved here is $0.0027(m_A/\sigma_A^4)$ which is regarded as optimum [16, 17]. The interaction between two ellipsoids with arbitrary orientations is assumed to be given by the Gay-Berne (GB)

*For correspondence: bbagchi@sscu.iisc.ernet.in

potential [18],

$$U_{GB} = 4\epsilon(\hat{r}, \hat{u}_1, \hat{u}_2) \left[\left(\frac{\sigma_0}{r - \sigma(\hat{r}, \hat{u}_1, \hat{u}_2) + \sigma_0} \right)^{12} - \left(\frac{\sigma_0}{r - \sigma(\hat{r}, \hat{u}_1, \hat{u}_2) + \sigma_0} \right)^6 \right] \quad (1)$$

where $\hat{u}_1, \hat{u}_2$ are the axial vectors of the ellipsoids 1 and 2. $\hat{r}$ is the vector along the intermolecular vector $r = r_2 - r_1$, where r_1 and r_2 denote the centers of mass of ellipsoids 1 and 2 respectively. $\sigma(\hat{r}, \hat{u}_1, \hat{u}_2)$ and $\epsilon(\hat{r}, \hat{u}_1, \hat{u}_2)$ are the orientation-dependent range and strength parameters respectively. σ and ϵ depend on the aspect ratio κ . The minor axis of the ellipsoid is equal to the diameter of the larger solvent and the major axis is 3 times that of the minor axis. Finally, the interaction between a sphere and an ellipsoid is accounted for by a modified GB-LJ potential given below

$$U_{Ei} = 4\epsilon_{Ei} \left[\left(\frac{\sigma(\theta)_{Ei}}{r} \right)^{12} - \left(\frac{\sigma(\theta)_{Ei}}{r} \right)^6 \right] \quad (2)$$

where 'E' denotes the ellipsoids and 'i' can be 'A' or 'B'. The expression for $\sigma(\theta)_{Ei}$ is available [19].

The ellipsoid in binary mixture system with the above mentioned potential is a well behaved system and it can also exhibit the experimentally observed anomalous viscosity dependence of the orientational correlation time [19]. Four ellipsoids were placed far from each other in a binary mixture of 500 particles with number of 'A' particles, $N_A = 400$ and number of 'B' type particles $N_B = 100$. The reduced temperature is expressed as, $T^* (= k_B T / \epsilon_A)$, the reduced pressure as, $P^* (= P \sigma_A^3 / \epsilon_{AA})$, and the reduced density as $\rho^* (= \rho \sigma_A^3)$. The time is scaled by $\tau = \sqrt{(m_A \sigma_A^2 / \epsilon_{AA})}$. The time step of the simulation is .002 τ and the system is equilibrated for 1.5×10^5 steps and the data collection step is 5×10^6 . The studies have been performed at $T^*=0.8$ and the $P^*=6$ and 10.

At $P^*=6$, both hopping and continuous motions exist in the system, thus the stress relaxation could not be directly correlated with the hopping. At $P^*=10$, only microscopic motion that survives is hopping. In a recent study, we have reported observation of correlated translational and orientational hopping [20] at this pressure. After extensive simulations we could find only two different kinds of motions. The translational hopping was either associated with correlated hopping of 5-6 nearest neighbors or it exhibited a motion in a ring like tunnel. While it is possible that other types of motion like isolated hopping can exist, we could not find them. The hopping rate was found to be 2×10^7 where both the type of motions occurred with almost equal frequency. In the following we focus on the stress relaxation and its relation vis-a-vis hopping at $P^*=10$ and $T^*=0.8$. The reduced density of the system is 1.41.

In figure 1 we show two different kinds of spatial hopping observed in our simulations. Both the hoppings




FIG. 1: (a) Shows the displacement of the 1st tagged ellipsoid over 1000 τ . There is a hopping of the ellipsoid around 300 τ . In the inset we plot the orientational correlation function obtained at different intervals. (b) Same as figure 1 (a) but for the 2nd tagged ellipsoid. The plots are at $P^*=10$ and $T^*=0.8$.

are associated with orientational hopping. In the inset we also plot the orientational time correlation functions (OCF), before, during and after the hopping. Figure 1(a) shows the trajectory of the first tagged ellipsoid and inset shows its orientational time correlation function, $\langle P_2(\hat{u}_i(0)\hat{u}_i(t)) \rangle / \langle P_2(\hat{u}_i(0)\hat{u}_i(0)) \rangle$, where P_2 is the second order Legendre polynomial. The hopping takes place in 20τ and the displacement is 0.5σ . Here the ellipsoid hopping is accompanied by hopping of 5-6 of its nearest neighbors. The OCF decays only during the period of hopping. In figure 1 (b) the trajectory of the second tagged ellipsoid and in the inset its orientational time correlation functions are shown. Note that in this case the displacement of the particle is large (1.1σ) and it also takes place over a very long period of time (50τ). Here we find that the tagged particle is moving in a ring like tunnel. Although the orientational correlation function decays during the hopping, its decay is less when compared to that of the 1st tagged particle.

The motions in a string like and ring like tunnel in a




FIG. 2: (a) The different components of the local stress auto-correlation (obtained from the sum of the stress of the 1st ellipsoid and its nearest neighbors) function obtained before hopping, between 320-330 τ . (b) is the same as 2 (a) but the components of the SAF are obtained after hopping between (340-350 τ). The plots are at $P^*=10$ and $T^*=0.8$.

deeply supercooled liquid have been reported earlier by other authors, although they did not discuss this large displacements over a long time for the ring like motion [21].

In the following part we discuss in details the local stress relaxation associated with these two different kinds of hoppings. The local stress around the ellipsoid is obtained by summing over the stress on the ellipsoid and its nearest neighbors. The stress has six components and the stress auto-correlation function is given by $\langle \sigma_{ij}(0)\sigma_{ij}(t) \rangle$, where $i,j=x,y,z$. In case of only Lennard-Jones fluid, $\sigma_{ij} = \sigma_{ji}$, but for particles interacting via GB and GB-LJ potential this is not so.

Figures 2 (a) and (b) show the SAF around the 1st tagged ellipsoid, before and after the period it is hopping, respectively. Before the hopping there is an anisotropy of the stress. The xy and yx components of the stress are much larger than the others and also their correlations cease to decay. *This anisotropy leads to a hopping of the ellipsoid, mostly in the z direction.* During the hopping there is a relaxation of the SAF in xy and yx components and after hopping all the components relax. Note that the total stress in all the components are also lower.

Figures 3 (a) and (b) show the SAF around the 2nd tagged ellipsoid, before and after the period it is hopping,




FIG. 3: (a) The different components of the local stress auto-correlation (obtained from the sum of the stress of the 2nd ellipsoid and its nearest neighbors) function obtained before hopping, between 1590-1630 τ . (b) is the same as 3 (a) but the components of the SAF are obtained after the hopping, between 1670-1710 τ . The plots are at $P^*=10$ and $T^*=0.8$.

respectively. Before the hopping there is an anisotropy of the stress. The yz and zy components of the stress are much larger than others, and also their correlations cease to decay. *This leads to the hopping of the ellipsoid, mostly in the x direction.* During and after the hopping there is an exchange of the stress. After the hopping although the yz component of the stress relaxes and also the $t=0$ value of all the components reduces, the SAF in the xz and zx components cease to decay. Thus, this kind of motion in a ring like tunnel does not lead to the relaxation of all the components of the stress.

The orientational correlation function of the 1st tagged particle relaxes more (inset of figure 1 (a)) compared to that of the 2nd tagged particle (inset of figure 1(b)) when computed in the respective intervals where they are hopping. From figure 2 and 3 we found that the local stress relaxation takes place when the 1st ellipsoid is hopping where as when the second ellipsoid is hopping although there is an exchange of stress between it's components, the local SAF does not completely relax. There is a direct connection between the local stress and the orientational relaxation functions, implying that rotation and viscosity are coupled even in a localized region.

In order to understand what happens to the surrounding of the local region when there is a relaxation of stress

due to hopping we have studied the stress auto correlation function of a bigger region of 2σ around the 1st ellipsoid. We found there are about 62-67 particles in this region. Although there is an anisotropy of the components of the stress in this larger region, this anisotropy cannot be correlated with the direction of hopping. Some of the components of SAF build up immediately after hopping and in a later time it relaxes. Similar analysis when done in a even bigger region shows that it takes longer for the stress of that region to relax and also the effect of the hopping is less.

In conclusion we demonstrated that the *direction* of the hopping of the tagged particle is determined by the anisotropy in the stress. Anisotropic stress relaxation is different when there is a many-particle hopping and there is a motion in a ring like tunnel. Although there is an exchange of stress between the components due to the particle motion, the stress relaxation is less in a ring like motion. Interestingly, the effect of hopping is found to spread over the adjoining region like ripples with the amplitude decreasing with increasing distance from the ellipsoid. We note that in the case of the second tagged el-

liploid (Fig.1b) although it translates more, the stress relaxation during its hopping is less. Thus suggesting that translational motion and viscosity are decoupled even in a localized region. On the other hand, the orientational relaxation and also the stress relaxation is more for the first ellipsoid. Thus suggesting that the orientational motion always remains coupled to viscosity. This is in agreement with the experimental results and in fact provides a microscopic explanation of the results which are known for a long time. There can be an apparent connection between the stress tensor and the momentum circulation. Thus it is possible that the non-decaying SAF implies that momentum circulation exists in a deeply super-cooled liquid. Since the anisotropic stress contributes to the free energy of the system, a change in the anisotropy drives the system from one free energy minimum to the other. When the anisotropy in the stress disappears and all the SAF relaxes then the system definitely moves to a lower free energy minimum.

This work was supported by a grant from CSIR, India. We thank Srikanth Sastry, Arnab Mukherjee and Rajesh Murarka for helpful discussions.

-
- [1] S. Sastry, P. G. Debenedetti and F. H. Stillinger, *Nature* **393**, 554 (1998).
 - [2] C. A. Angell, *Chem. Rev.* **90**, 523 (1990); Luz-Maria Martinez and C. A. Angell, (private communication)
 - [3] A. Mukherjee, S. Bhattacharyya, and B. Bagchi, **116**, 4577 (2001).
 - [4] L. Angelani, G. Parisi, G. Ruocco and G. Vilianni, *Phys. Rev. Lett.* **81**, 4648 (1998).
 - [5] C. D. Michele and D. Leporini, *Phys. Rev. E* **63**, 036701 (2001); *Phys. Rev. E* **63**, 036702 (2001).
 - [6] G. Hinze, D. D. Brace, S. D. Gottke, and M. D. Fayer, *Phys. Rev. Lett.* **84**, 2437 (2000); *J. Chem. Phys.* **113**, 3723 (2000).
 - [7] G. P. Johari and M. Goldstein, *J. Chem. Phys.* **53**, 2372 (1970); *ibid*, **55**, 4245 (1971).
 - [8] M.D. Ediger, C. A. Angell and S. R. Nagel, *J. Phys. Chem.* **100**; M. T. Cicerone and M. D. Ediger, *J. Chem. Phys.* **104**, 7210, **103**, 5684 (1996); G. Heuberger and H. Sillescu, *J. Chem. Phys.* **100**, 15255 (1996).
 - [9] L. A. Deschenes and D. A. Vanden Bout, *Science* **292**, (2001).
 - [10] C. K. Majumdar, *Solid State Com.* **9**, 1087 (1971).
 - [11] P. C. Hohenberg and B. I. Halperin, *Rev. Mod. Phys.* **49**, 435 (1977); L. D. Landau and E. M. Lifshitz, *Theory of Elasticity* (Pergamon Press, U. K 1981).
 - [12] S. Sastry, *Phys. Rev. Lett.* **85**, 590 (2000); S. Sastry, *Nature*, **409**, 164 (2001).
 - [13] W. Kob and H. C. Andersen, *Phys. Rev. E*, **51**, 4626 (1995); W. Kob and H. C. Andersen, *Phys. Rev. Lett.*, **73**, 1376 (1994).
 - [14] K. Vollmayr, W. Kob and K. Binder, *J. Chem. Phys.* **105**, 4714 (1996); F. Sciortino, W. Kob and P. Tartaglia, *Phys. Rev. Lett.* **83**, 3214 (1999); B. Coluzzi, G. Parisi and P. Verrocchio, *J. Chem. Phys.* **112**, 2933 (2000).
 - [15] H.C. Andersen et al. Rapport d'activite scientifique du CECAM, pp 82-115 [7.4, 7.4.3].
 - [16] D. Brown and J. H. R. Clarke, *Mol. Phys.* **51**, 1243 (1984).
 - [17] J. M. Haile and H. W. Graben, *J. Chem. Phys.* **73**, 2412 (1980).
 - [18] J. G. Gay and B. J. Berne, *J. Chem. Phys.*, **74**, 3316 (1981).
 - [19] S. Bhattacharyya and B. Bagchi, *J. Chem. Phys.*, **115**, 9065 (2001).
 - [20] S. Bhattacharyya, A. Mukherjee and B. Bagchi, *J. Chem. Phys.*, **117**, 2741 (2002).
 - [21] C. Bennemann, C. Donati, J. Baschnagel, and S. C. Glotzer, *Nature (London)* **399**, 246 (1999).